

# CLICK PREMIUM

GROUP	5	HERBICIDE
GROUP	27	HERBICIDE

A suspension concentrate formulation containing 330 g/l Terbutylazine and 70g/l Mesotrione. A selective herbicide for the control of annual grasses and broad-leaved weeds in forage and grain maize.

**24 HOUR EMERGENCY NUMBER: +44(0)1763 212100. In the event of an emergency, call the National Poisons Centre, Beaumont Hospital at 01 809 2566 or 01 837 9964.**

## SAFETY INFORMATION FOR PROFESSIONAL USE ONLY

### WARNING



- **Harmful if swallowed**
- **Suspected of damaging the unborn child**
- **May cause damage to organs through prolonged or repeated exposure**
- **Very toxic to aquatic organisms, may cause long term adverse effects in the aquatic environment**

- Do not handle until all safety precautions have been read and understood.
- Do not breathe spray.
- Wear protective gloves/protective clothing.
- Wash skin thoroughly after handling.
- IF exposed or concerned: Get medical advice/attention.
- Collect spillage.
- Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty triple rinsed clean containers which can be disposed of as non-hazardous waste.

PCS Number 07269

UFI: JF06-PT17-Q10Y-SY05

Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.

**To avoid risks to human health and the environment comply with the instructions for use.**

**SIPCAM UK LTD CONDITIONS OF SUPPLY** - All goods supplied by us are of high grade, but as we cannot exercise control over their mixing or use, all conditions or warranty, statutory or otherwise, as to the quality or fitness of our goods for any purpose are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

#### Authorisation Holder

SIPCAM OXON S.P.A.  
Via Sempione 195,  
20016 Pero (MI) Italy

#### Marketing Company

SIPCAM UK Ltd, 4C Archway House,  
The Lanterns, Melbourn Street,  
Royston, Herts SG8 7BX

Tel +44(0)1763 212100  
Fax +44(0)1763 212101

Net contents:

# 5Le



Scan to view the Safety Data Sheet

Alternatively, download the Safety Data Sheet from [sipcamuk.co.uk](http://sipcamuk.co.uk) or contact your supplier.

[sipcamuk.co.uk](http://sipcamuk.co.uk)

  
**SIPCAM**  
UK

# FOR PROFESSIONAL USE ONLY

## FOR USE ONLY AS AN AGRICULTURAL HERBICIDE

Crops	Maximum individual dose: (L product/hectare)	Maximum total dose: (L product/hectare)	Maximum number of treatments	Latest time of application
Maize (forage and grain)	1.5 L/ha	1.5L ha per crop	–	Up to 8 leaves unfolded (GS18)
Other restriction: Product may only be used from 2 leaves unfolded stage (GS12). Do not apply on sands or very light soils. Residual weed control will be reduced on soils with over 10% organic matter. Use shall be limited to one application of any terbuthylazine containing product, every three years on the same field.				
<b>READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.</b>				

## SAFETY PRECAUTIONS

### Operator protection:

Wear suitable protective gloves when handling concentrate.  
For use by tractor mounted/trailed sprayers.

### Environmental protection:

To protect aquatic organisms, respect an unsprayed buffer zone of at least 5m to surface water bodies.

To protect non-target plants, respect an unsprayed buffer zone of 5m to non-agricultural land unless using 90% drift reducing nozzles.

Do not contaminate water with the product or its container.

Do not clean application equipment near surface water.

Avoid contamination via drains from farmyards and roads.

### Storage and disposal:

Keep in original container, tightly closed in a safe place.

Rinse container thoroughly by using an integrated pressure device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

Do not re-use container for any purpose.

## DIRECTIONS FOR USE

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

CLICK PREMIUM is a herbicide for the control of annual grass and broad-leaved weeds in forage and grain maize.

### Restrictions/Warnings

For best results on open ground crops, young weeds are most susceptible.

Do not use on seed crops or on sweetcorn varieties.

CLICK PREMIUM should be applied to actively growing weeds, post-emergence.

Treatment in poor growing conditions or on dry soil may give less reliable control.

Do not spray when foliage is wet.

Under adverse conditions mild to moderate chlorosis may be observed on sprayed leaves, this effect is usually transient and does not affect yield.

Take extreme care to avoid drift onto plants outside the treated area.

Ensure spray swaths do not overlap.

Addition of an adjuvant or tank mix wetting agent is not recommended.

### Weed Control

Post-emergence use only - Dose rate depends on weed type at maize crop growth stage GS 12-16.

Annual broad leaved weeds	Dose rate L/ha	Weed growth stage
Fat Hen	1.0	Emergence to start of branching (GS20)
Common Chickweed	1.0	Emergence to start of flowering (GS25-30)
Black Nightshade	1.0	Emergence to 6 leaves
Field pansy	1.0	Emergence to 10 leaves
Annual Grasses	Dose rate L/ha	Weed growth stage
Annual meadow grass *	1.5	Emergence to mid tiller

\*moderate control

### Mixing

Never prepare more spray solution than is required.

Half fill the tank with clean water and start the agitation. To ensure thorough mixing of the product, invert the container several times before opening. Add the required quantity of CLICK PREMIUM to the spray tank while re-circulating. Fill up the tank with water and continue agitation until spraying is completed.

On emptying the container, rinse container thoroughly by using an integrated pressure rinsing device or manually

rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

Continue agitation during spraying.

### Application

Ensure good, even spray cover of the target crop and weeds. Apply using medium quality spray through a conventional field crop sprayer. Do not leave spray liquid in the sprayer for longer periods (i.e. overnight)

Apply CLICK PREMIUM in 200-300L of water/ha. Higher volume may be required for dense or well developed weed populations.

### Sprayer cleaning

After spraying, thoroughly clean and flush out application machinery with a minimum of three rinses, to ensure that all traces of product are removed.

Disposal of spray tank washings should be in accordance with local legislation. Unused product and containers must not be allowed to enter water and sewage systems.

## FOLLOWING CROPS AND RECULTIVATION

### Recultivation

Ploughing is recommended prior to reseeded. Slight crop effects seen after emergence are usually transitory. Maize can be reseeded in case of crop failure.

### Crop rotation

#### Autumn

Winter wheat (including durum wheat), winter barley, and rye grass can follow a maize crop. Deep ploughing (greater than 15cm) is necessary before drilling oilseed rape

#### Spring

Forage and grain maize, rye grass, spring wheat and spring barley may be sown in spring following use of CLICK PREMIUM but cannot be treated with terbuthylazine containing products.

DO NOT SOW spinach, sugar and fodder beet, peas, beans lettuce or cabbage in the year following application.